

## **PHALLUS ATROVOLVATUS, A NEW SPECIES FROM COSTA RICA**

by

F.D. CALONGE<sup>1</sup>, H. KREISEL<sup>2</sup> & M. MATA<sup>3</sup>

<sup>1</sup>Real Jardín Botánico, CSIC, Plaza de Murillo 2., E-28014 Madrid, Spain.

e-mail: calonge@ma-rjb.csic.es

<sup>2</sup>Zur Schwedenschanze 4, D-17498 Potthagen, Germany.

e-mail: Hanns.Kreisel@gmx.de

<sup>3</sup>INBio, Instituto Nacional de Biodiversidad, A. P. 22-3100, Santo Domingo, Heredia, Costa Rica.

e-mail: mmata@inbio.ac.cr

**Summary.** F.D. CALONGE, KREISEL H; & M. MATA (2005). *Phallus atrovolvatus*, a new species from Costa Rica. *Bol. Soc. Micol. Madrid* 29: 5-8.

*Phallus atrovolvatus* is described as a new species from Costa Rica, which belongs to Section *Clautriavia* (Pat.) Kreisel. This taxon is characterized by showing a white indusium, rugulose to merulioid receptaculum (pileus) surface, beige greenish gleba and a black volva. Notes on its taxonomy and ecology are also added.

**Key words:** *Basidiomycotina*, *Phallales*, *Phallus*, taxonomy, ecology, Costa Rica, Central America.

**Resumen.** F.D. CALONGE, KREISEL H; & M. MATA (2005). *Phallus atrovolvatus*, especie nueva encontrada en Costa Rica. *Bol. Soc. Micol. Madrid* 29: 5-8.

Se propone *Phallus atrovolvatus* como especie nueva, el cual fue colectado en Costa Rica y pertenece a la Sección *Clautriavia* (Pat.) Kreisel. El nuevo taxon se caracteriza por su indusio blanco, receptáculo con superficie rugulosa a merulioide, gleba beige con tonos verdosos y volva negra. Se incluyen notas sobre su taxonomía y ecología.

**Palabras clave:** *Basidiomycotina*, *Phallales*, *Phallus*, taxonomía, ecología, Costa Rica, América Central.

### **INTRODUCTION**

Regarding the taxonomy of *Phallus*, numerous have been the publications on this genus in the world (VENTENAT, 1798; FISCHER, 1888-1905; MÖLLER, 1895; LLOYD, 1909; BOEDIJN, 1932; PILAT, 1958; DRING & RAYNER, 1967; DRING & ROSE, 1976; LIU, 1984; MILLER & MILLER, 1988; ZANG & PETERSEN, 1989; etc.); however most of them appear fragmentary and dispersed, being necessary and urgent the confection of a world monograph. In

1996 KREISEL published a preliminary survey of the genus *Phallus*, but without including any systematic key. As a matter of fact there is a great need to have a work which contains such a tool for all the interested, both amateurs and professionals, on this genus.

One of us (FDC), had the opportunity to spend six months in Costa Rica, looking forward to the study of the *Gasteromycetes* and *Pezizales* of that country. As a consequence of that stay several contributions have been carried out since then (CALONGE & MATA, 2002; CALONGE &



Fig. 1.- *Phallus atrovolvatus*. A colony with several basidiomata where it is possible to observe unopen "eggs", and well-developed basidiomata showing receptacle, indusium, pseudostipe and a black volva. Holotypus.

CARRANZA, 2003; CALONGE & *al.*, 2003; CALONGE & *al.*, 2005), and we had the chance to collect the material of a rare *Phallus*, which has resulted being a new taxon.

The terminology used here follows that published by CALONGE (1998).

## DESCRIPTION

***Phallus atrovolvatus*** Kreisel & Calonge sp. nov. Figs. 1-2.

Etym.: *atrovolvatus*, refers to the black colour of the volva.

*Ovum* 20-30 mm diametrum, nigrum. *Basidioma* mature cum pseudostipite cylindrico, 40-100 x 10-20 mm, album, spongioso. *Indusium* album. *Receptaculo* trunco-conico cum apice perforato, 10-20 mm alto et 15-30 mm diametro ad basim, ruguloso-merulioideo. *Gleba* cremea viridis cum odore suave, non foetido.

*Volva nigra. Sporae ellipsoideae*, 3-5 x 1-1.5  $\mu$ m, laeves, chlorini-hyalinae. *Gregarious ad residuis lignum putridum*.

COSTA RICA: Limón, La Amistad Caribe, Cahuita, gregarious on wood chips, among grasses, 12-VII-2001, leg. F. D. Calonge. **Holotypus**: INB3814309-A. **Isotypus**: MA-Fungi 59261.

Unexpanded basidioma globose to ovoid, 20-30 mm diam., black (Fig. 1). Exoperidium membranous, endoperidium gelatinous, hyaline. At maturity a cylindric pseudostipe develops up to 40-100 x 10-20 mm, white, spongy (Figs. 1-2). Receptacle conical-truncate with a rugulose to meruloid surface, perforate apex (Fig. 2), 10-20 mm high and 15-30 mm wide at the base. Indusium expanded to midway between receptacle and volva, with lacerate margin, white to pale yellow (Fig. 2), 30-50 mm long, consisting of a latticed, pseudoparenchymatous pendent membrane (Figs. 1-2) attached to the apex beneath the



receptacle. Gleba beige to yellowish green (Figs. 1-2), without any repellent odour, somewhat sweet, aromatic. Spores ellipsoid,  $3\text{--}5 \times 1\text{--}1.5 \mu\text{m}$ , smooth, hyaline with a weak yellowish tint. Volva black, paler later (Figs. 1-2). It grows gregariously on wood chips and plant debris, among grasses.

## DISCUSSION

The effort dedicated to the study of these fungi in different parts of the world, during the last years, have brought the publication of six new species: *Phallus taipeiensis* Liu & Bau (LIU & BAU, 1982), *P. echinovolvatus* (M. Zang, D. R. Zheng & Z. X. Hu) Kreisel (ZANG & *al.*, 1988), *P. yunnanensis* (M. Zang & R. H. Petersen) Kreisel (ZANG & PETERSEN, 1989), *Phallus nanchangensis* Z. Z. He (HE, 1989) in China; *Phallus minusculus* Kreisel & Calonge (CALONGE & KREISEL, 2002) in tropical Africa and *Phallus pygmaeus* Baseia (BASEIA & *al.*, 2003) in Brazil.

On the other hand, there have been synonymized

other species, such as *P. rugulosus* (E. Fisch.) O. Kuntze, *P. novae-hollandiae* Corda, *P. canariensis* Mont. All three under *P. rubicundus* (Bosc) Fr. Another group: *P. moelleri* Lloyd, *P. callichrous* (A. Möller) Lloyd have been placed under *P. indusiatus* Vent.: Pers. (CUNNINGHAM, 1944).

In other instances the holotypus of several species seems to be missing, such as *P. subtilis* (A. Möller) Lloyd, *P. costatus* (Penz.) Lloyd, *P. tenuis* (E. Fisch.) O. Kunze, *P. favosus* (Penz.) E. Fisch., *P. lauterbachii* (Henn.) Kreisel, *P. callichrous* (A. Möller) Lloyd, etc.

*Phallus atrovolvatus* seems to behave as a rare species, since it has never been found again after its discovery, being close to *P. merulinus* (Berk.) Lloyd, which differs in having a white volva and growth on soil. The re-examination of 29 collections of *Phallus* belonging to the USJ, CR and INB herbaria, trying to find more samples of *P. atrovolvatus*, has resulted in 26 collections corresponding to *P. indusiatus* and three to *P. duplicatus*.



Fig. 2.- *Phallus atrovolvatus*. Basidiomata showing all the characters described in the previous figure, but with a paler volva in the lied down one. Holotypus

Thus, regarding our material the combination of characters such as a well-developed white indusium; receptacle with rugulose to meruloid surface; gleba beige with a sweet, non repellent odour; white pseudostipe; lack of any rose or purple tints on the indusium, and a black volva award to *Phallus atrovolutus* a unique identity. That is why we propose it as a new species.

## ACKNOWLEDGEMENTS

One of us (FDC) wishes to express his gratitude to the Spanish Ministry of Education and Science, for financial support during his stay in Costa Rica. We are grateful to the staff of INBio for their help during the preparation of this work, and to Dr. B. Liu for his kind sending of information.

## BIBLIOGRAPHY

- BASFIA, I.G., T.B. GIBERTONI & I.C. MAIA (2003). *Phallus pygmaeus*, a new minute species from tropical rain forest. *Mycotaxon* 85: 77-79.
- BOEDJUN, K.B. (1932). The Phallineae of the Netherlands East Indies. *Bull. Jard. Bot. Buitenzorg* III, 12: 71-103.
- CALONGE, F.D. (1998). *Gasteromycetes* I *Lycoperdales*. *Nidulariales*. *Phallales*. *Sclerodermatales*. *Tulostomatales*. *Fl. Mycol. Iberica* 3: 1-271.
- CALONGE, F.D. & H. KREISEL (2002). *Phallus minusculus* sp. nov. from Tropical Africa. *Feddes Repertorium* 113: 600-602.
- CALONGE, F.D. & M. MAIA (2002). *Plectania carranzae* sp. nov. (*Ascomycotina*) from Costa Rica. *Mycotaxon* 81: 237-241.
- CALONGE, F.D. & J. CARRANZA (2003). *Tulostoma matae* sp. nov. (*Gasteromycetes*), found in Costa Rica. *Bol. Soc. Micol. Madrid* 27: 37-42.
- CALONGE, F.D.; T. ITURRIAGA; M. MATA & J. CARRANZA (2003). Contribución al estudio de los *Pezizales* (*Ascomycotina*) de Costa Rica. *Bol. Soc. Micol. Madrid* 27: 21-32.
- CALONGE, F.D.; M. MAIA & J. CARRANZA (2005). Contribución al catálogo de los *Gasteromycetes* de Costa Rica. *An. Real Jardín Bot. Madrid* 62:23-45.
- CUNNINGHAM, G.H. (1944). *The Gasteromycetes of Australia and New Zealand*. J. Cramer, Vaduz (reprint 1979).
- DRING, D.M. & R.W. RAYNER (1967). Some *Gasteromycetes* from Eastern Africa. *J. East. Afr. Nat. Hist. Soc.* 26: 5-46.
- DRING, D.M. & A.C. ROSE (1976). Additions to West African phalloid fungi. *Kew Bull.* 31: 741- 751
- FISCHER, E. (1888-1905). *Phalloideae*. In Saccardo, P. A., *Syll. Fungorum* 7: 3-14 and 469; 9: 262-264; 11: 152-154; 14: 254; 16: 224-228; 17: 212. Patavii.
- HE, Z.Z. (1989). A new species of the genus *Phallus* from China. *Acta Mycol. Sinica* 8: 98-100.
- KREISEL, H. (1996). A preliminary survey of the genus *Phallus* sensu lato. *Czech Mycol.* 48: 273-281
- LIU, B. (1984). The *Gasteromycetes* of China. *Nova Hedwigia*, Beih., 76: 1-235.
- LIU, B. & Y. S. BAO (1982) The genus *Phallus* from China. *J. Shanxi Univ.* 4: 95-102.
- LLOYD, C.G. (1909). Synopsis of the known phalloids. *Bull. Lloyd Library* 13: 1-96.
- MILLER, O.K. & H.H. MILLER (1988). *Gasteromycetes, morphological and development features with keys to orders, families and genera*. Ed. Mad River Press, Eureka.
- MÖLLER, A. (1895). *Brasilische Pilzblumen*. Ed. G. Fischer, Jena.
- PILÁT, A. (1958). *Phallus*. In Pilát (ed.), *Fl. CSR. Gasteromycetes*. : 58-71.
- VENTENAT, E.P. (1798). Dissertation sur le genre *Phallus*. *Mem. Inst. Nat. Sci. Arts Sc. Math. Phys.* 1: 503- 523.
- ZANG, M. & R.H. PETERSEN (1989). *Endophallus*, a new genus in the *Phallaceae* from China. *Mycologia* 81: 486- 489.
- ZANG, M., D.R. ZENG & Z.X. HU (1988). A new species of the genus *Dictyophora* from China. *Mycotaxon* 33: 145-148.